

LATTICE STUDY OF NUCLEON PROPERTIES WITH DOMAIN WALL FERMIONS

SHOICHI SASAKI

*Department of Physics, University of Tokyo,
7-3-1 Hongo, Tokyo 113-0033, Japan
E-mail: ssasaki@phys.s.u-tokyo.ac.jp*

Domain wall fermions (DWF) are a new fermion discretization scheme with greatly improved chiral symmetry. Our final goal is to study the nucleon spin structure through lattice simulation using DWF. In this paper, we present our current progress on two topics toward this goal: 1) the mass spectrum of the nucleon excited states and 2) the iso-vector vector and axial charges, g_V and g_A , of the nucleon.

1. Introduction

The RIKEN-BNL-Columbia-KEK QCD Collaboration has been pursuing the domain-wall fermion (DWF) method¹ in lattice quantum chromodynamics (QCD). In DWF an extra fifth dimension is added to the lattice. By manipulating the *domain-wall* structure of the fermion mass in this fifth dimension, we control the number of light fermion species in the other four space-time dimensions. These light fermions possess exact chiral symmetry in the limit of an infinite fifth dimension. In particular: 1) fermion near-zero mode effects are well understood², 2) explicit chiral symmetry breaking induced by a finite extra dimension is described by a single residual mass parameter, which is very small in the present calculation, in the low-energy effective lagrangian², and 3) non-perturbative renormalization works well³.

DWF is a promising new approach for treating fermions on the lattice. However, we need several tests of DWF in the baryon sector to reach our final goal of establishing the spin structure of the nucleon from first principles. Here we report our recent studies of the mass spectrum of the nucleon and its excited states and the nucleon matrix elements of the iso-vector vector and axial charges in quenched lattice QCD with DWF. Although most of the latter results are preliminary, the conclusive results in the former subject have been reported in Ref.[4].

2. Nucleon excited states

First, we discuss the mass spectrum of the nucleon N and its excited states (the negative-parity nucleon N^* and the positive-parity first excited nucleon N') by means of a systematic investigation utilizing two distinct interpolating

operators $B_1^\pm$ and $B_2^\pm$. For an explanation of those operators, see Ref.[4]. Our quenched DWF calculation was employed on lattice with size $16^3 \times 32 \times 16$, gauge coupling $\beta = 6/g^2 = 6.0$, and domain wall height $M_5 = 1.8$. Additional details of our simulation can be found in Ref.[4].

In Fig.1 we show the low-lying nucleon spectrum as a function of the quark mass, m_f in lattice units ($a^{-1} \approx 1.9$ GeV set from $aM_\rho=0.400(8)$ in the chiral limit). B_1^+ gives the ground-state nucleon mass N (cross). The N^* mass estimates (square and diamond) are extracted from both B_1^- and B_2^- . The corresponding experimental values for N and N^* are marked with lower and upper stars. Both N^* mass estimates extracted from two distinct operators agree with each other. The large N - N^* mass splitting is clearly evident.

In contrast to the negative parity operators, we find that the mass estimates from a second operator, B_2^+ , are considerably larger than the ground state obtained from B_1^+ . This suggests that B_2^+ has negligible overlap with the nucleon ground state and provides a signal for the positive-parity *excited* nucleon N' . To justify this possibility, we employ a sophisticated approach which utilizes the transfer matrix of a 2×2 correlation function⁵ constructed from both B_1^+ and B_2^+ . The diagonalization of the transfer matrix yields the excited state⁵. Fig.2 shows a comparison of the fitted mass from $\langle\langle B_2^+(t)\bar{B}_2^+(0)\rangle\rangle$ (circle) and the estimated mass from the average effective mass given by the smaller eigenvalue of the transfer matrix (bullet). The cross symbol corresponds to the nucleon ground state mass evaluated from the larger eigenvalue of the transfer matrix, which is quite consistent with the fitted mass from $\langle\langle B_1^+(t)\bar{B}_1^+(0)\rangle\rangle$.

Another important conclusion can be drawn from Fig.1 and Fig.2. In the heavy quark mass region, the ordering of the negative-parity nucleon (N^*) and the positive-parity excited nucleon (N') is inverted relative to experiment. This remarkable result was originally reported in our early paper⁴ and subsequently confirmed in Ref.[6]. Further systematic calculation is required to determine whether this ordering switches to the observed ordering as one approaches the chiral limit.

3. Nucleon matrix elements

The nucleon (iso-vector) axial charge g_A is a particularly interesting quantity. We know precisely the experimental value $g_A = 1.2670(35)$ from neutron beta decay. Why does g_A deviate from unity in contrast to vector charge, $g_V = 1$? The simple explanation is given by the fact that the axial current is only partially conserved in the strong interaction while the vector current is exactly conserved. Thus, the calculation of g_A is an especially relevant test of

the chiral properties of DWF in the baryon sector. In addition, calculation of g_A is an important first step in studying polarized nucleon structure functions since $g_A = \Delta u - \Delta d$ where $\langle p, s | \bar{q}_f \gamma_5 \gamma_\mu q_f | p, s \rangle = 2s_\mu \Delta q_f$ with $s^2 = -1$ and $s \cdot p = 0$.

We follow the standard practice^{8,9} for the calculation of g_V and g_A . The three-point functions for the local vector current $J_V^f = \bar{q}_f \gamma_A q_f$ and axial current $J_A^f = \bar{q}_f \gamma_5 \gamma_i q_f$ ($i = 1, 2, 3$) are defined by

$$G_\Gamma^{u,d}(t, t') = \text{Tr}[P_\Gamma \sum_{\vec{x}, \vec{x}'} \langle T B_1(\vec{x}, t) J_\Gamma^{u,d}(\vec{x}', t') \bar{B}_1(0, 0) \rangle] \quad (1)$$

with $\Gamma = V$ (vector) or A (axial) where $P_V = P_+ = (1 + \gamma_4)/2$ and $P_A = P_+ \gamma_i \gamma_5$. For the axial current, the three-point function is averaged over $i = 1, 2, 3$. The lattice estimates of vector and axial charges can be derived from the ratio between two- and three-point functions

$$g_\Gamma^{\text{lattice}} = \frac{G_\Gamma^u(t, t') - G_\Gamma^d(t, t')}{G_N(t)}, \quad (2)$$

where $G_N(t) = \text{Tr}[P_+ \sum_{\vec{x}} \langle T B_1(\vec{x}, t) \bar{B}_1(0, 0) \rangle]$.

Recall that in general lattice operators $\mathcal{O}_{\text{lat}}$ and continuum operator $\mathcal{O}_{\text{con}}$ are regularized in different schemes. The operators are related by a renormalization factor Z_O : $\mathcal{O}_{\text{con}}(\mu) = Z_O(a\mu) \mathcal{O}_{\text{lat}}(a)$. This implies that the continuum value of vector and axial charges are given by $g_\Gamma = Z_\Gamma g_\Gamma^{\text{lattice}}$. In the case of conventional Wilson fermions, the renormalization factor Z_A is usually estimated in perturbation theory (Z_A differs from unity because of explicit symmetry breaking). For DWF, the conserved axial current receives no renormalization. This is not true for the lattice local current. An important advantage with DWF, however, is that the lattice renormalizations, Z_V and Z_A , of the local currents are the same³ so that the ratio $(g_A/g_V)^{\text{lattice}}$ directly yields the continuum value g_A .

Our preliminary results are analyzed on 200 quenched gauge configurations at $\beta = 6.0$ on a $16^3 \times 32 \times 16$ lattice with $M_5 = 1.8^7$. We choose a fixed separation in time of the nucleon interpolating operators, $t = t_{\text{source}} - t_{\text{sink}}$ and $t' < t$, with currents inserted in between. We take $t_{\text{source}} = 5$ and $t_{\text{sink}} = 21^7$.

In Fig.3 we show the dependence of the vector renormalization, $Z_V = 1/g_V^{\text{lattice}}$ on the location of current insertions. A good plateau is observed so that Z_V is certainly well behaved. The value 0.763(5) at $m_f = 0.02$ (obtained by averaging over time slices denoted by the solid line in Fig.3) agrees well with $Z_A = 0.7555(3)$, which was obtained from a completely different calculation of meson two-point correlation functions based on the relation $\langle A_\mu^{\text{conserved}}(t) \bar{q} \gamma_5 q(0) \rangle = Z_A \langle A_\mu^{\text{local}}(t) \bar{q} \gamma_5 q(0) \rangle^2$.

For the axial charge, g_A^{lattice} , plateaus are seen for $10 \leq t \leq 16$ in Fig.4, so the charge ratios $(g_A/g_V)^{\text{lattice}}$ at each m_f are averaged in this time slice range. We find that there is a fairly strong dependence of $(g_A/g_V)^{\text{lattice}}$ on m_f as be shown in Fig.5. A simple linear extrapolation to $m_f = 0$ yields $g_A = 0.62(13)$, which is roughly 1/2 the experimental value⁷. However, a simple linear ansatz may not describe the data in which case the result in the chiral limit may be even smaller.

To compare to results using Wilson fermions, we plot g_A versus the square of the π - ρ mass ratio in Fig.6. Our result is given by the bullets. We also include two heavier mass points (with large errors) from an earlier simulation using a larger separation between the source and the sink. The triangles and diamonds are from Wilson simulations at relatively strong⁸ and weak coupling⁹. At first glance, the DWF and Wilson fermions seem to be in rough agreement except for the lightest point. However, our DWF results have a strong mass dependence. This may be a finite volume effect; the Wilson results at strong coupling were simulated on a lattice with a volume which is twice ours.

A couple of comments on the mass dependence of our data are in order. First, it is interesting to note that our results appear consistent with the value 5/3 (marked as star) in the heavy quark limit, while the others seem inconsistent with this limit. Second, our results may also be consistent with vanishing in the chiral limit. This can be explained through the axial Ward-Takahashi identity which governs g_A . If the PCAC relation $m_\pi^2 \propto m_f$ is modified, for example by chiral logarithms, the nucleon matrix element of the pseudoscalar density does not develop a pole as $m_f \rightarrow 0$, and the r.h.s of the identity vanishes in the chiral limit. Thus, g_A must also vanish as $m_f \rightarrow 0$. Indeed, we already know that the PCAC relation for the pion mass is modified in the quenched approximation by two effects: zero-modes of the Dirac operator and the quenched chiral logarithm². Further investigation of the Ward-Takahashi identity is under way.

4. Conclusions

We have explored several nucleon properties in quenched lattice QCD using domain wall fermions toward our final goal of studying the spin structure of the nucleon from first principles.

Our quenched DWF calculation reproduces very well the large mass splitting between the nucleon $N(939)$ and its parity partner $N^*(1535)$ ⁴. We have also calculated the mass of the first positive-parity excited state $N'(1440)$ by the diagonalization of a 2×2 matrix correlator and confirmed that it is heavier than the negative-parity excited state $N^*(1535)$ ⁴. A remaining puzzle is

whether or not a switching of N^* and N' occurs close to the chiral limit.

A preliminary calculation of iso-vector vector and axial charges shows that all the relevant three-point functions are well behaved. Z_V determined from the nucleon matrix element of the vector current agrees closely with that from an NPR study of quark bilinears³ and a direct calculation using meson correlation functions². This indicates that $g_V = 1$ and $Z_V = Z_A$ are mutually satisfied in our quenched DWF calculation. However, a linear extrapolation of g_A to the chiral limit yields a value which is a factor of two smaller than the experimental value. We are currently investigating the Ward-Takahashi identity that governs g_A to shed light on this behavior. We also plan to check related systematic effects arising from finite lattice volume and quenching (for example quenched chiral logarithms, zero modes, and the absence of the full pion cloud), especially in the lighter quark mass region.

Acknowledgments

The author would like to thank the organizers of NSTAR2001 for an invitation. This work was done in collaboration with Tom Blum and Shigemi Ohta as a part of the RIKEN-BNL-Columbia-KEK QCD Collaboration. We thank RIKEN, Brookhaven National Laboratory, and the U.S. Department of Energy for providing the facilities essential for the completion of this work.

References

1. An early review of domain wall fermions is given in T. Blum, Nucl. Phys. **B** (Proc. Suppl.) **73**, 167 (1999) ; For a recent review see P. Vranas, Nucl. Phys. **B** (Proc. Suppl.) **94** 177 (2001).
2. T. Blum, *et al.*, hep-lat/0007038.
3. T. Blum, *et al.*, hep-lat/0102005.
4. S. Sasaki, T. Blum and S. Ohta, hep-lat/0102010; S. Sasaki, Nucl. Phys. **B** (Proc. Suppl.) **83**, 206 (2000) and hep-ph/0004252.
5. M. Lüscher and U. Wolff, Nucl. Phys. **B339**, 222 (1990).
6. F.X. Lee, hep-lat/0011060; D. Richards, hep-lat/0011025.
7. T. Blum, S. Ohta and S. Sasaki, Nucl. Phys. **B** (Proc. Suppl.) **94**, 295 (2001).
8. M. Fukugita *et al.*, Phys. Rev. Lett. **75**, 2092 (1995).
9. M. Göckeler *et al.*, Phys. Rev. **D53**, 2317 (1996).




Figure 1. N and N^* (square and diamond) masses versus the quark mass m_f in lattice units. Note the large N - N^* mass splitting which is within 10% (in the chiral limit) of the experimental value (bursts).




Figure 2. The mass of the positive-parity excited state (circles) is too high compared to the nucleon ground state (cross) to account for the observed splitting.




Figure 3. Dependence of vector renormalization, $Z_V = 1/g_V^{\text{lattice}}$, on t' , at $m_f = 0.02$. A good plateau is observed.




Figure 4. The lattice axial charge, g_A^{lattice} , at $m_f = 0.02$. A good plateau is seen in the range $10 \leq t \leq 16$.




Figure 5. Dependence of $(g_A/g_V)^{\text{lattice}}$ on m_f .




Figure 6. g_A versus $(m_\pi/m_\rho)^2$.